

Some stable vector bundles with reducible theta divisors

Arnaud BEAUVILLE

Introduction

Let C be a curve of genus g , and E a vector bundle on C , of rank r ; assume that the *slope* $\mu := \frac{1}{r} \deg E$ of E is an integer. Let J^ν be the variety (isomorphic to the Jacobian of C) parametrizing line bundles of degree $\nu := g - 1 - \mu$ on C . We say that E *admits a theta divisor* if $H^0(E \otimes L) = 0$ for L general in J^ν . If this is the case, the locus

$$\Theta_E = \{L \in J^\nu \mid H^0(E \otimes L) \neq 0\}$$

has a natural structure of effective divisor in J^ν , the *theta divisor* of E . Its class in $H^2(J^\nu, \mathbf{Z})$ is $r\theta$, where $\theta \in H^2(J^\nu, \mathbf{Z})$ is the class of the principal polarization. This (generalized) theta divisor plays a key role in the recent work on vector bundles on curves – see for instance [B] for an overview.

If E admits a theta divisor, it is semi-stable (otherwise E contains a sub-bundle F of slope $> \mu$, and by Riemann-Roch $H^0(F \otimes L) \neq 0$, and therefore $H^0(E \otimes L) \neq 0$, is non-zero for all $L \in J^\nu$). The converse does not hold, at least in rank ≥ 4 : Raynaud has constructed examples of stable vector bundles with no theta divisor [R]. Further examples have been constructed recently by Popa [P].

If E is semi-stable but not stable, its theta divisor (if it exists) is not integral: more precisely, E admits a filtration with stable quotients $E_1, \dots, E_p$, and we have $\Theta_E = \Theta_{E_1} + \dots + \Theta_{E_p}$. One may ask, conversely, if the reducibility of Θ_E implies that E is not stable. A counter-example has been given by Raynaud (unpublished), who constructed a rank 2 stable vector bundle on a curve of genus 3 with reducible theta divisor. Such an example can only occur on a special curve, because in rank 2 the divisor Θ_E characterizes the vector bundle E [B-V], and on a general curve the only reducible divisors on J^ν with cohomology class 2θ are the theta divisors of rank 2 decomposable bundles.

We describe in this note a counter-example of a different nature, namely a family of stable vector bundles of rank g which exist on any curve of genus g . They are defined by the exact sequence

$$0 \rightarrow M_L \longrightarrow H^0(C, L) \otimes_{\mathbf{C}} \mathcal{O}_C \xrightarrow{\text{ev}_L} L \rightarrow 0$$

where L is a line bundle generated by its global sections, and ev_L the evaluation map. These vector bundles have been intensively studied, notably by Green and

Lazarsfeld (see in particular [L]), Paranjape and Ramanan [P-R], and more recently in [P] and [F-M-P]. In the latter paper the authors determine the theta divisor of M_K and of its exterior powers; we will do the same here in the case of a line bundle L of degree $2g$ (so that M_L has rank g). We will prove, in a somewhat more precise form, the following result:

Theorem.— *Let C be a non-hyperelliptic curve, and L a sufficiently general line bundle of degree $2g$ on C . The vector bundle M_L and its exterior powers $\Lambda^2 M_L, \dots, \Lambda^{g-1} M_L$ are stable and admit a reducible theta divisor.*

An interesting extra feature of our examples is that there exists a semi-stable, decomposable vector bundle on C with the same theta divisor as M_L ; thus in rank ≥ 3 the divisor Θ_E does not characterize the bundle E any more.

Notation

We fix a curve C of genus g ; except in Remark 2 below, we assume throughout that C is not hyperelliptic. For $d \in \mathbf{Z}$, we denote by J^d the translate of the Jacobian of C parametrizing line bundles of degree d on C , and by C_d the locus of effective divisor classes in J^d . If $p, q \in \mathbf{Z}$ the difference variety $C_p - C_q$ lies in J^{p-q} .

I. The Theta divisor of E_L

Let L be a line bundle of degree $2g$ on the curve C . It is spanned by its global sections, so we have an exact sequence

$$0 \rightarrow M_L \rightarrow H^0(L) \otimes_{\mathbf{C}} \mathcal{O}_C \rightarrow L \rightarrow 0,$$

where M_L is a rank g vector bundle. We put $E_L := M_L^*$.

Though this will not be used in the sequel, let us recall the geometric interpretation of E_L . Let φ be the morphism of C into the projective space $\mathbf{P} := \mathbf{P}(H^0(L))$ defined by the linear system $|L|$; in view of the Euler exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbf{P}} \rightarrow H^0(L)^* \otimes_{\mathbf{C}} \mathcal{O}_{\mathbf{P}}(1) \rightarrow T_{\mathbf{P}} \rightarrow 0,$$

we have $E_L = \varphi^* T_{\mathbf{P}} \otimes L^{-1}$.

The vector bundle E_L has rank g and determinant L , hence slope 2.

Proposition 1.— a) *The vector bundle E_L has a theta divisor*

$$\Theta_{E_L} = (C_{g-2} - C) + \Theta_{L \otimes K^{-1}} \quad \text{in } J^{g-3}.$$

b) *E_L is semi-stable; it is stable if and only if L is very ample.*

Proof : We will first compute set-theoretically the theta divisor Θ_{M_L} of M_L . By definition this is the set of line bundles $P \in J^{g+1}$ such that the multiplication map $\mu : H^0(L) \otimes H^0(P) \rightarrow H^0(L \otimes P)$ is not injective. Let us distinguish three cases:

- (i) If $h^0(P) > 2$ we have $\dim H^0(L) \otimes H^0(P) > \dim H^0(L \otimes P)$, thus $P \in \Theta_{M_L}$.
- (ii) Assume that $h^0(P) = 2$ and that the pencil $|P|$ has a base point. Both spaces $H^0(L) \otimes H^0(P)$ and $H^0(L \otimes P)$ have the same dimension $2g + 2$. If μ is injective, it is surjective, and the linear system $|L \otimes P|$ has a base point; this is impossible since $\deg(L \otimes P) = 3g + 1$. Thus we have again $P \in \Theta_{M_L}$.
- (iii) Finally assume that $|P|$ is a base-point free pencil. From the exact sequence

$$0 \rightarrow P^{-1} \longrightarrow H^0(P) \otimes_{\mathbf{C}} \mathcal{O}_C \longrightarrow P \rightarrow 0$$

we get an exact sequence

$$0 \rightarrow H^0(L \otimes P^{-1}) \longrightarrow H^0(L) \otimes_{\mathbf{C}} H^0(P) \xrightarrow{\mu} H^0(L \otimes P) ;$$

thus μ is not injective in that case if and only if $H^0(L \otimes P^{-1}) \neq 0$.

The line bundles P in case (i) and (ii) are exactly those which can be written $P'(x)$, for some point x of C and some line bundle P' in J^g with $h^0(P') \geq 2$; the ones in case (iii) are those of the form $L \otimes P'^{-1}$, with $P' \in \Theta \subset J^{g-1}$. Since Θ_{E_L} is the image of Θ_{M_L} by the isomorphism $L \mapsto K \otimes L^{-1}$ of J^{g+1} onto J^{g-3} , we obtain (still set-theoretically) $\Theta_{E_L} = (C_{g-2} - C) \cup \Theta_{L \otimes K^{-1}}$. Now $C_{g-2} - C$ is an irreducible divisor with cohomology class $(g-1)\theta$ (see e.g. [F-M-P], Prop. 3.7), and $\Theta_{L \otimes K^{-1}}$ is a (ordinary) theta divisor; since Θ_{E_L} has cohomology class $g\theta$, we get the equality a).

Since E_L admits a theta divisor, it is semi-stable. Moreover, its stable components are $L' := L \otimes K^{-1}$ and a rank $(g-1)$ bundle. If E_L is not stable, L' is either a sub- or a quotient bundle of E_L . The latter case cannot occur since E_L is generated by its global sections and L' is not. Now using the exact sequence

$$0 \rightarrow L^{-1} \otimes L'^{-1} \longrightarrow H^0(L)^* \otimes_{\mathbf{C}} L'^{-1} \longrightarrow E_L \otimes L'^{-1} \rightarrow 0$$

and Serre duality we see that $\text{Hom}(L', E_L)$ is zero if and only if the multiplication map $H^0(L) \otimes H^0(L) \rightarrow H^0(L^2)$ is surjective, that is, L is normally generated [G-L]. By [G-L], Thm. 1, this is the case if and only if L is very ample. ■

Remarks.— 1) If L is not very ample, we have $L = K(D)$, with D an effective

divisor of degree 2. The snake lemma applied to the commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & M_K & \longrightarrow & H^0(K) \otimes \mathcal{O}_C & \longrightarrow & K \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & M_L & \longrightarrow & H^0(L) \otimes \mathcal{O}_C & \longrightarrow & L \longrightarrow 0
\end{array}$$

provides an exact sequence $0 \rightarrow M_K \rightarrow M_L \rightarrow \mathcal{O}_C(-D) \rightarrow 0$; thus E_L is an extension of E_K by $\mathcal{O}_C(D)$. This extension is non-trivial, as we already observed that $\mathcal{O}_C(D)$ cannot be a quotient of E_L .

2) If C is hyperelliptic, the divisor $C_{g-2} - C$ is equal to Θ_H , where H is the hyperelliptic pencil on C . By specialization we get $\Theta_{E_L} = (g-1)\Theta_H + \Theta_{L \otimes K^{-1}}$. The line bundle L is not linearly normal [L-M], so E_L is not stable.

The difference variety $C_{g-2} - C$ is the theta divisor of the bundle E_K [P-R]; therefore:

Corollary 1.— *Assume that L is very ample. The stable bundle E_L and the decomposable bundle $E_K \oplus (L \otimes K^{-1})$ have the same theta divisor. ■*

The equality still holds of course when L is not very ample, but becomes immediate, since in that case the second bundle is the sum of the stable components of the first one.

In view of the results of [B-N-R], this corollary can be rephrased as follows. Let $\mathcal{SU}_C(g)$ be the moduli space of semi-stable rank g vector bundles on C with trivial determinant, and let $\mathcal{L}$ be the positive generator of $\text{Pic}(\mathcal{SU}_C(g))$ (the *determinant bundle*). Let $B_{\mathcal{L}}$ be the base locus of the linear system $|\mathcal{L}|$.

Corollary 2.— *The map $\varphi_{\mathcal{L}} : \mathcal{SU}_C(g) - B_{\mathcal{L}} \rightarrow \mathbf{P}(H^0(\mathcal{L}))$ defined by the line bundle $\mathcal{L}$ is not injective.*

Indeed this map can be identified with the map which associates to a vector bundle its theta divisor [B-N-R]. Twisting E_L and $E_K \oplus (L \otimes K^{-1})$ by a line bundle λ on C with $\lambda^{-g} = L$, we get two different points of $\mathcal{SU}_C(g) - B_{\mathcal{L}}$ with the same image under $\varphi_{\mathcal{L}}$. ■

II. The Theta divisor of $\Lambda^p E_L$

We now consider the exterior power $\Lambda^p E_L$; this is a vector bundle of rank $\binom{g}{p}$ and slope $2p$, so its theta divisor, if it exists, lie in J^{g-1-2p} .

Proposition 2.— *Let $1 \leq p \leq g-1$. If L is general enough, the vector bundle $\Lambda^p E_L$ is stable and admits a theta divisor¹*

$$\Theta_{\Lambda^p E_L} = (C_{g-p-1} - C_p) + (C_{g-p} - C_{p-1} + K \otimes L^{-1}) .$$

Proof: We first prove that $\Lambda^p E_L$ admits a theta divisor when L is general enough. Since this is an open property, it is sufficient to prove this for a particular choice of L : we take $L = K(D)$, with D an effective divisor of degree 2. The exact sequence

$$0 \rightarrow \mathcal{O}_C(D) \rightarrow E_L \rightarrow E_K \rightarrow 0$$

(Remark 1) gives rise to an exact sequence

$$0 \rightarrow \Lambda^{p-1} E_K(D) \rightarrow \Lambda^p E_L \rightarrow \Lambda^p E_K \rightarrow 0 .$$

Since by [F-M-P] each exterior power $\Lambda^q E_K$ admits a theta divisor, so does $\Lambda^p E_L$.

Let us now prove that the theta divisor Θ_{E_L} , when it exists, is given by the formula of the Proposition. The divisor $C_q - C_{g-1-q}$ has cohomology class $\binom{g-1}{q}\theta$ ([F-M-P], Prop. 3.7), so both sides of the formula have cohomology class $\binom{g}{p}\theta$. It suffices therefore to prove that each component of the right hand side is contained in $\Theta_{\Lambda^p E_L}$.

As in [P] and [F-M-P], we will use the following observation of Lazarsfeld [L]: if $x_1, \dots, x_{g-1}$ are generic points of C , there is an exact sequence

$$0 \rightarrow \bigoplus_{i=1}^{g-1} \mathcal{O}_C(x_i) \longrightarrow E_L \longrightarrow L(-\sum x_i) \rightarrow 0$$

which gives rise as above to an exact sequence

$$0 \rightarrow \bigoplus_{i_1 < \dots < i_p} \mathcal{O}_C(x_{i_1} + \dots + x_{i_p}) \longrightarrow \Lambda^p E_L \longrightarrow \bigoplus_{j_1 < \dots < j_{g-p}} L(-x_{j_1} - \dots - x_{j_{g-p}}) \rightarrow 0 .$$

This gives:

- $H^0(\Lambda^p E_L(-x_1 - \dots - x_p)) \neq 0$, hence the inclusion $C_{g-p-1} - C_p \subset \Theta_{\Lambda^p E_L}$;
- $H^0(\Lambda^p M_L \otimes L(-x_1 - \dots - x_{g-p})) \neq 0$, hence $H^0(\Lambda^p M_L \otimes L(-D)) \neq 0$ for all D in $C_{g-p} - C_{p-1}$; by Serre duality this gives $H^0(\Lambda^p E_L \otimes K \otimes L^{-1}(D)) \neq 0$, hence the inclusion $C_{g-p} - C_{p-1} + K \otimes L^{-1} \subset \Theta_{\Lambda^p E_L}$.

It remains to prove that $\Lambda^p E_L$ is stable. Since L is generic, E_L is stable (Proposition 1), so $\Lambda^p E_L$ is *polystable* – that is, direct sum of stable bundles with the same

¹ The second term is the translate of $C_{g-p} - C_{p-1} \subset J^{g+1-2p}$ by the element $K \otimes L^{-1}$ of J^{-2} .

slope $2p$. If $\Lambda^p E_L$ is not stable for L generic, it is decomposable for all values of L ; we will see that this is not the case when L is of the form $K(D)$, with D effective of degree 2. In that case we have by Remark 1 an exact sequence

$$0 \rightarrow \Lambda^{p-1} E_K(D) \longrightarrow \Lambda^p E_L \longrightarrow \Lambda^p E_K \rightarrow 0$$

where $\Lambda^{p-1} E_K(D)$ and $\Lambda^p E_K$ are stable with slope $2p$; if $\Lambda^p E_L$ is decomposable, this exact sequence splits. The following easy lemma shows that this is not the case, and thus concludes the proof of the Proposition. ■

Lemma.— *Let X be a scheme over a field of characteristic 0, and let*

$$(\mathcal{E}) \quad 0 \rightarrow M \rightarrow E \rightarrow F \rightarrow 0$$

be a non-split exact sequence of vector bundles on X , with $\text{rk } M = 1$. The associated exact sequences

$$(\Lambda^p \mathcal{E}) \quad 0 \rightarrow \Lambda^{p-1} F \otimes M \longrightarrow \Lambda^p E \longrightarrow \Lambda^p F \rightarrow 0$$

do not split for $1 \leq p \leq \text{rk } F$.

Proof: Let $i : F^* \otimes M \rightarrow \mathcal{H}om(\Lambda^p F, \Lambda^{p-1} F \otimes M)$ be the linear map deduced from the interior product. A straightforward computation shows that the class of the extension $(\Lambda^p \mathcal{E})$ in $H^1(X, \mathcal{H}om(\Lambda^p F, \Lambda^{p-1} F \otimes M))$ is the image by $H^1(i)$ of the class of the extension $(\mathcal{E})$ in $H^1(X, F^* \otimes M)$. But in characteristic zero i admits a retraction $c^{-1} \rho$, where $c = \binom{\text{rk } F - 1}{p-1}$ and $\rho : \mathcal{H}om(\Lambda^p F, \Lambda^{p-1} F \otimes M) \rightarrow F^* \otimes M$ is the map deduced from the interior product $\Lambda^p F^* \otimes \Lambda^{p-1} F \rightarrow F^*$. Thus $H^1(i)$ is injective, and the lemma follows. ■

As in section I this gives:

Corollary 1.— *The vector bundles $\Lambda^p E_L$ and $\Lambda^p E_K \oplus (\Lambda^{p-1} E_K \otimes L \otimes K^{-1})$ have the same theta divisor. In particular, if L is general enough, the map $\varphi_{\mathcal{L}} : \mathcal{SU}_C(\binom{g}{p}) - B_{\mathcal{L}} \longrightarrow \mathbf{P}(H^0(\mathcal{L}))$ defined by the line bundle $\mathcal{L}$ is not injective.*

Let us conclude by a link with the main theme of [F-M-P], the so-called *minimal resolution conjecture* for the curve C embedded into $\mathbf{P}^g := \mathbf{P}(H^0(L))$. We have to refer to [F-M-P] for the statement of the conjecture, which is a bit technical. Let us just say that it describes, for all general finite subsets $\Gamma \subset C$ of cardinality $\geq g + 1$, the minimal graded resolution of the ideal I_{Γ} of Γ in the coordinate ring $S = \mathbf{C}[X_0, \dots, X_g]$ of $\mathbf{P}^g$. By Corollary 1.8 of [F-M-P], this conjecture holds if and only if each of the bundles $\Lambda^p E_L$ admit a theta divisor. Thus:

Corollary 2.— *The curve C , embedded into $\mathbf{P}^g$ by a general linear system of degree $2g$, satisfies the “minimal resolution conjecture” in the sense of [F-M-P]. ■*

REFERENCES

- [B] A. BEAUVILLE: *Vector bundles on curves and generalized theta functions: recent results and open problems*. Current topics in complex algebraic geometry, 17–33, Math. Sci. Res. Inst. Publ. **28**, Cambridge Univ. Press (1995).
- [B-N-R] A. BEAUVILLE, M.S. NARASIMHAN, S. RAMANAN: *Spectral curves and the generalised theta divisor*. J. Reine Angew. Math. **398** (1989), 169–179.
- [B-V] S. BRIVIO, A. VERRA: *The theta divisor of $SU_C(2, 2d)^s$ is very ample if C is not hyperelliptic*. Duke Math. J. **82** (1996), 503–552.
- [F-M-P] G. FARKAS, M. MUSTĂŢĂ, M. POPA: *Divisors on $\mathcal{M}_{g,g+1}$ and the Minimal Resolution Conjecture for points on canonical curves*. Preprint math.AG/0104187, to appear in Duke Math. J.
- [G-L] M. GREEN, R. LAZARSEFELD: *On the projective normality of complete linear series on an algebraic curve*. Invent. Math. **83** (1986), 73–90.
- [L] R. LAZARSEFELD: *A sampling of vector bundle techniques in the study of linear series*. Lectures on Riemann Surfaces, World Scientific Press, Singapore (1989), pp. 500–559.
- [L-M] H. LANGE, G. MARTENS: *Normal generation and presentation of line bundles of low degree on curves*. J. Reine Angew. Math. **356** (1985), 1–18.
- [P] M. POPA: *On the base locus of the generalized theta divisor*. C. R. Acad. Sci. Paris **329** (1999), Série I, 507–512.
- [P-R] K. PARANJAPE, S. RAMANAN: *On the canonical ring of a curve*. Algebraic geometry and commutative algebra, vol. II, 503–516, Kinokuniya, Tokyo (1988).
- [R] M. RAYNAUD: *Sections des fibrés vectoriels sur une courbe*. Bull. Soc. Math. France **110** (1982), 103–125.

Arnaud BEAUVILLE
 Institut Universitaire de France
 &
 Laboratoire J.-A. Dieudonné
 UMR 6621 du CNRS
 UNIVERSITÉ DE NICE
 Parc Valrose
 F-06108 NICE Cedex 02